arXiv:1110.4288v1 [cond-mat.dis-nn] 19 Oct 2011

Ordinary Percolation with Discontinuous Transitions

Stefan Boettcher and Vijay Singh
Department of Physics, Emory University, Atlanta, GA 30322; USA

Robert M. Ziff
*Center for the Study of Complex Systems and Department of Chemical Engineering,
University of Michigan, Ann Arbor, MI 48109-2136; USA*

Percolation on finitely ramified lattices, those where the loss of a finite number of bonds can forestall end-to-end transport when the number of nodes N increases, is generally considered to be trivial[1]. They percolate only at full bond density, $p = 1$. Examples are the one-dimensional lattice and fractals such as the Sierpinski gasket[2]. We show here that by dressing up such lattices with small-world bonds[3], in a way that the resulting network becomes infinitely ramified, an entirely novel percolation transition at a critical point $p_c < 1$ emerges. Approaching this transition from below, many clusters of already diverging, yet, sub-extensive, sizes combine *explosively*[4] at p_c such that the resulting cluster becomes extensive. That is, the usual order parameter P_∞, describing the probability of any node to be part of the largest cluster, jumps instantly to a finite value at p_c. Simple examples of this transition are provided by small-world networks consisting of a one-dimensional lattice combined with various hierarchies of long-range bonds[5] that reveal many features of the transition explicitly.

The percolation properties of networks are of significant interest; without percolation, any large-scale transport or communication through the network ceases. Much research has been dedicated to the understanding of percolation on randomly grown, complex networks[6]. Yet, the engineering of artificial networks with well-controlled features seems desirable. Indeed, there has been considerable interest in the properties of spatial networks, linking real-world geometry with small-world effects[7,8]. In particular, networks possessing hyperbolic and/or hierarchical features[5,7,9–14] relate to actual transport systems such as for air travel, routers, and social interactions. Hierarchical networks exhibit a number of novel features, as shown with *exact* methods below, most notably a discontinuous transition in the formation of an extensive percolating cluster for ordinary, random bond addition. For random networks, such an "explosive" percolation transition, one that requires *correlated* bond additions, has been proposed by Achlioptas et al.[4] only recently. Intense research is ongoing into the dynamics of correlated cluster formation[15–22], with the existence of a discontinuity still in doubt[23–26]. Understanding of explosive cluster formation could be significantly advanced when the discontinuity can be studied rigorously, merely by adding bonds randomly on an engineered network.

A simple example of a hierarchical network is developed in Fig.1(a). In its recursive construction, bonds from the preceding generation are built up in the next generation according to a fixed pattern up to a generation $k \to \infty$. Applying a venerable technique called real-space renormalization group (RG)[27] to the complete network in effect *reverses* the recursive built-up of the network, as shown in Fig.1(b). Starting with a network of size $N = 2^k+1$, the probability of two neighboring sites to be directly linked is $T_0 = p$. From that, we can recursively determine the probability T_n, $0 \le n \le k$, that two sites originally separated by 2^n positions in the horizontal direction have a direct path between them. Finally, $T_{n=k}$ provides the probability for *any* end-to-end path in the network; for the two end-points all paths are direct. As explained in Fig.1(b), these probabilities satisfy the recursion

$$T_{n+1} = p + (1-p)T_n^2, \qquad (T_0 = p)\,. \tag{1}$$

For infinitely large networks we find $\lim_{k\to\infty} T_{n=k} = T^*$ from the stationary points, e.g., Eq. (1) yields the fixed point $T^* \equiv 1$ for all p and a line of fixed points,

$$T^*(p) = \frac{p}{1-p} \qquad \left(p \le \frac{1}{2}\right), \tag{2}$$

that intersect at $p = \frac{1}{2}$. Analysis shows[5] that Eq. (2) is a stable fixed point for all $0 < p < \frac{1}{2}$ which describes the asymptotic behavior of large networks with a finite probability $0 < T^*(p) < 1$ to possess an end-to-end path. For $\frac{1}{2} \le p \le 1$, there is always such a path, i.e., $T^* = 1$.

It is remarkable that percolation on these networks has a number of features entirely absent from basic lattices[5,12,14]. Specifically, on a lattice (like the 1d line) the existence of an end-to-end path is synonymous with the existence of an extensive cluster, giving each node a finite probability $P_\infty(p)$ to be connected to it. This is not generally true for these hyperbolic-shaped networks. Typically, there are *two* transitions: below the lower transition p_l there are only finite clusters and no end-to-end paths, and above the upper transition $p_u > p_l$ there is an extensive cluster and a definite end-to-end path, as on any lattice. However, between p_l and p_u exists a regime with diverging but sub-extensive clusters and a variable probability for an end-to-end path, as given by Eq. (2). Those sub-extensive clusters scale with a new fractal exponent $0 \le \Psi(p) < 1$.[30] Accordingly, the order parameter P_∞[1,27] is strictly non-zero only above p_u, making it the thermodynamically correct transition



Figure 1: (a) Pattern for the recursive generation of a hierarchical network (displayed at full bond density, $p = 1$). In $k = 0$-th generation, the network consists of a single bond between two end-nodes (open circles). In successive generations $k = 1, 2, \ldots$, each (horizontal) lattice bond is replaced by two of its kind, with a new (black) node in-between, and a long-range bond (shaded arcs). In generation k the network has $N = 2^k + 1$ nodes and $L = 2^{k+1} - 1$ bonds for a (sparse) average degree $2L/N \sim 4$ for large k. (b) The n-th RG step reduces each three-node graph-let into a single lattice bond of renormalized percolation probability for a direct path between nodes T_{n+1}, which derives from that probability, T_n, of the prior bonds and the probability p to have a long-range bond. T_{n+1} percolates if either the long-range bond exists (probability p, irrespective of T_n) or both lattice bonds are present (probability $(1-p)T_n^2$), leading to Eq. (1). Reversing the pattern set in (a), the RG starts at $n = 0$ from an unrenormalized network of generation k with uniform probabilities p for the long-range bonds as well as for the direct path to neighboring nodes, $T_0 = p$.



Figure 2: (a) Schematic for generating functions $T_n(x)$ and $S_n(x,y)$. The open circles represent the end-nodes, shaded areas indicate clusters that either span (T_n) or do not span (S_n) between the end-nodes. Clusters that do not reach an end-node are ignored. (b) All diagrams contributing to $T_{n+1}(x)$ or $S_{n+1}(x,y)$ in the nth RG step. The remaining end-nodes (always-open circles) are not counted in the generating functions; the (black) connecting nodes increment the cluster size and are taken into account by multiplying x or y. The contribution of each graph-let is: (i) $xpT_n^2(x)$, (ii) $xpT_n(x)S_n(x,x)$, (iii) $xpT_n(x)S_n(x,x)$, (iv) $pS_n(x,1)S_n(1,x)$, (v) $x(1-p)T_n^2(x)$, (vi) $x(1-p)T_n(x)S_n(x,y)$, (vii) $y(1-p)T_n(y)S_n(x,y)$, and (viii) $(1-p)S_n(x,1)S_n(1,y)$. As tallied up in Eqs. (5,6), graph-lets (i)-(v) span end-to-end and contribute to $T_{n+1}(x)$, while (vi)-(viii) do not span and contribute to $S_{n+1}(x,y)$.

point, $p_c = p_u$. In the present network it is $p_l = 0$; a more elaborate network[5,28] with non-trivial p_l and p_u is mentioned below.

To reveal the nature of the transition, we obtain the exact P_∞ from cluster generating functions. To this end, we introduce two basic quantities: the probability $t_i^{(n)}(p)$ that both end-nodes are connected to the same cluster of size i, and the probability $s_{i,j}^{(n)}(p)$ that the left (right) end-node is connected to a cluster of size i (j) but these clusters are not the same. For convenience the end-nodes are *not* counted in the cluster size, as shown in Fig.2(a). The corresponding generating functions are defined as

$$T_n(x) = \sum_{i=0}^{\infty} t_i^{(n)}(p)x^i, \tag{3}$$

$$S_n(x,y) = \sum_{i=0}^{\infty}\sum_{j=0}^{\infty} s_{i,j}^{(n)}(p)x^i y^j. \tag{4}$$

The recursion relations for these generating functions



Figure 3: Plot of $P_\infty(p)$, evaluated after $n=10^1-10^5$ iterations of the recursions in Eq. (10), evolving into a discontinuity at $p\to p_c=\frac{1}{2}$ with $P_\infty(p_c)=0.609793\ldots$. Convergence is slowest just below p_c, since finite-size corrections decay slowly with $N^{\Psi(p)-1}$ with $1-\Psi(p)\sim(p_c-p)/\ln 2$ for $p\to p_c$ from Eq. (13).

can be obtained by considering all possible triangular graph-lets, like in Fig. 1(b), but now also taking cluster sizes into account, as shown in Fig.2(b). For each graph-let, the type of connectivity between each of the nodes is checked, see Fig.2(a), and the bonds are assigned a value $T_n(x)$ or $S_n(x,y)$, depending on the type of cluster each represents. As in Fig. 1(b), small world bonds are merely assigned the probability p or $1-p$ for being present or not. Marking the increment in cluster size, the inner node provides a factor of x or y for its adjacent cluster, or unity if it remains isolated. The contribution of each graph-let to the next generation is the product of the weights of the three bonds and the intermediate node; all eight of these are determined in Fig.2(b). These provide the recursions

$$\begin{aligned} T_{n+1}(x) &= xT_n^2(x)+p\left[2xT_n(x)S_n(x,x)\right.\\ &\quad\left.+S_n(x,1)S_n(1,x)\right], \qquad (5)\\ S_{n+1}(x,y) &= (1-p)\left[xT_n(x)S_n(x,y)+\right.\\ &\quad\left.+yT_n(y)S_n(x,y)+S_n(x,1)S_n(1,y)\right] \qquad (6)\end{aligned}$$

initiated with $T_0(x)=p$, $S_0(x,y)=1-p$. Note that both equations reduce to Eq. (1) for $x=y=1$, where Eqs. (3,4) give $T_n(1)=1-S_n(1,1)=T_n$.

It is convenient to rewrite the recursions in Eqs. (5,6) in terms of functions of a single variable x only. To this end, we define the functions $\Sigma_n(x)\equiv S_n(x,x)$ and $\mathcal{S}_n(x)\equiv S_n(x,1)=S_n(1,x)$ that, combined into a vector $\vec{V}=(T,\Sigma,\mathcal{S})$, lead to

$$\vec{V}_{n+1}(x)=\vec{\mathcal{F}}\left(\vec{V}_n(x),x\right) \qquad (7)$$

with a function-vector $\vec{\mathcal{F}}$ of non-linear components

$$\vec{\mathcal{F}}\left(\vec{V},x\right) = \begin{pmatrix} xT^2+2xpT\Sigma+p\mathcal{S}^2\\ (1-p)\left(2xT\Sigma+\mathcal{S}^2\right)\\ (1-p)(1+xT)\mathcal{S}\end{pmatrix}. \qquad (8)$$

The mean size $\langle s_{\max}\rangle_k$ of the (likely largest) cluster connected to the end-nodes and, hence, the order parameter $P_\infty(p)=\lim_{k\to\infty}\langle s_{\max}\rangle_k/N_k$ result from the first moment of the generating functions via the first derivative in x at $x=1$,

$$\langle s_{\max}\rangle_k=T_k'+\Sigma_k'\sim N_k^{\Psi(p)}, \qquad (9)$$

for a network of size $N_k=2^k+1\to\infty$. Expanding Eq. (7) to first order in $\epsilon\equiv1-x\to0$ yields to zeroth order Eq. (7) evaluated at $x=1$, such that each component simply reproduces Eq. (1) again. To order ϵ, we find an inhomogeneous linear recursion for $\vec{V}_n'=\vec{V}_n'(x=1)$

$$\vec{V}_{n+1}' = \vec{V}_n'\circ\frac{\partial\vec{\mathcal{F}}}{\partial\vec{V}}\left(\vec{V}_n,1\right)+\frac{\partial\vec{\mathcal{F}}}{\partial x}\left(\vec{V}_n,1\right) \qquad (10)$$

with the Jacobian matrix

$$\frac{\partial\vec{\mathcal{F}}}{\partial\vec{V}}\left(\vec{V},1\right)=\begin{bmatrix} 2T+2p\Sigma, & 2pT, & 2p\mathcal{S}\\ 2(1-p)\Sigma, & 2(1-p)T, & 2(1-p)\mathcal{S}\\ (1-p)\mathcal{S}, & 0, & (1-p)(1+T)\end{bmatrix} \qquad (11)$$

and inhomogeneity

$$\frac{\partial\vec{\mathcal{F}}}{\partial x}\left(\vec{V},1\right) = \begin{pmatrix} T^2+2pT\Sigma\\ 2(1-p)T\Sigma\\ (1-p)T\mathcal{S}\end{pmatrix} \qquad (12)$$

which depend on n through $\vec{V}_n(x=1)=(T_n,1-T_n,1-T_n)$. Since each network at $n=0$ only consists of end-nodes, all clusters are empty and $\vec{V}_0'=\vec{0}$.

For large n, i.e., near the fixed point $\vec{V}^*=(T^*,1-T^*,1-T^*)$, it is easy to show that the inhomogeneity in Eq. (10) is subdominant, leaving a simple, homogeneous, linear system. For each component of $\vec{V}_n'$, the dominant contribution then arises from the eigenstate with the largest eigenvalue of $\frac{\partial\vec{\mathcal{F}}}{\partial\vec{V}}\left(\vec{V}^*,1\right)$. Below the transition, $0\leq p<p_c=\frac{1}{2}$, we use Eq. (2) for T^* to obtain as the largest eigenvalue $\lambda=1+2p$, such that $\langle s_{\max}\rangle_k\sim\lambda^k$ for $k\to\infty$. Then Eq. (9) yields the fractal exponent

$$\Psi(p)=\frac{\ln\lambda}{\ln2}=\frac{\ln(1+2p)}{\ln2} \qquad (13)$$

for sub-extensive cluster sizes, with $0\leq\Psi<1$, implying $P_\infty\equiv0$ here.

Above and *at* the transition, $p_c\leq p\leq1$, $T^*=1$ and Eq. (11) provides uniformly $\lambda=2$ as the largest eigenvalue (i.e., $\Psi\equiv1$), indicating the existence of an extensive cluster. But unlike for a continuous transition, where $P_\infty(p)\to0$ for $p\to p_c^+$, Eq. (10) provides for a monotone increasing probability $\langle s_{\max}\rangle_k/N_k$ for any node to be connected to a spanning cluster, even at $p=p_c=\frac{1}{2}$, as shown in Fig. 3. In fact, the continuity of $P_\infty(p)$ is interrupted merely because $T^*\equiv1$ becomes an *unstable* fixed point of Eq. (1) below p_c. There the stable



Figure 4: Depiction of the non-planar Hanoi network HNNP of finite size[5]. Starting from a $1d$-lattice (horizontal lines), a set of long-range bonds (arced lines) is added hierarchically that break planarity. RG on this network remains exact. Nodes on the lowest two levels of the hierarchy have degree 3, then degree 5, 7, etc., comprising a fraction of 1/2, 1/4, 1/8, etc., of all nodes, resulting in an average degree 4.

branch transitions to Eq. (2) that lacks extensive clusters. Hence, it is the intersection of two stable branches of fixed points T^* that causes a discontinuous transition.

We believe, that such a transition is *generic* for this type of hierarchical network. For instance, on the non-planar Hanoi network HNNP[5] described in Fig. 4, an equivalent but far more elaborate calculation results in a $p_l = 0.319445181\ldots$ as the root of an octic polynomial in p, while $p_u = \left(3-\sqrt{5}\right)/2 = 0.381966\ldots$. We find $\Psi \equiv 0$ below p_l, at which the exponent Ψ jumps to a finite value, followed by a domain with $\ln 1.66639/\ln 2 \approx 0.736726 \leq \Psi\left(p\right) < 1$ for $p_l \leq p < p_u$. At p_u, again, two lines of fixed points intersect, resulting in an extensive cluster, $\Psi \equiv 1$, above and at $p_u = p_c$, causing a discontinuity in P_∞. For $p \to p_c$, finite-size corrections decay as $N^{\Psi(p)-1}$, where $1-\Psi\left(p\right) \sim \left(p_c - p\right)^\kappa$ with a $\kappa > 1$, signaling subtle variations in the cluster formation dynamics between networks.

We have shown that a hierarchy of small-world bonds grafted onto a finitely ramified lattice results generically in an explosive percolation transition. This differs significantly from other models, where a non-local set of bonds or a global growth parameter is considered to achieve explosive percolation[4,15–26]. At this point, the precise conditions to be imposed on the hierarchy of long-range bonds for obtaining this transition are not entirely clear. However, they have to result in an infinitely ramified network[1] to provide $p_c < 1$, as the example of the Hanoi network HN3 demonstrates[5]. As well, a finitely ramified lattice seems essential, as studies of hierarchical networks with small-world bonds imposed on infinitely ramified 2d-lattices[12,14] appear to result in infinite-order transitions, which have been observed in many other networks[29].

---

[1] D. Stauffer and A. Aharony, *Introduction to Percolation Theory,* 2^{nd} *Ed.* (CRC Press, Boca Raton, 1994).

[2] H. Taitelbaum, S. Havlin, P. Grassberger, and U. Moening, J. Phys. A: Math. Gen. **23**, 371 (1990).

[3] D. J. Watts and S. H. Strogatz, Nature **393**, 440 (1998).

[4] D. Achlioptas, R. M. D'Souza, and J. Spencer, Science **323**, 1453 (2009).

[5] S. Boettcher, J. L. Cook, and R. M. Ziff, Phys. Rev. E **80**, 041115 (2009).

[6] A.-L. Barabasi, *Linked: How Everything Is Connected to Everything Else and What It Means for Business, Science, and Everyday Life* (Plume Books, 2003).

[7] M. Boguñá, D. Krioukov, and K. C. Claffy, Nature Physics **5**, 74 (2009).

[8] M. Barthelemy, Physics Reports **499**, 1 (2011).

[9] A. Trusina, S. Maslov, P. Minnhagen, and K. Sneppen, Phys. Rev. Lett. **92**, 178702 (2004).

[10] J. S. Andrade, H. J. Herrmann, R. F. S. Andrade, and L. R. da Silva, Phys. Rev. Lett. **94**, 018702 (2005).

[11] M. Hinczewski and A. N. Berker, Phys. Rev. E **73**, 066126 (2006).

[12] A. N. Berker, M. Hinczewski, and R. R. Netz, Phys. Rev. E **80**, 041118 (2009).

[13] D. Krioukov, F. Papadopoulos, M. Kitsak, A. Vahdat, and M. Boguñá, Phys. Rev. E **82**, 036106 (2010).

[14] T. Hasegawa, M. Sato, and K. Nemoto, Phys. Rev. E **82**, 046101 (2010).

[15] N. A. M. Araujo and H. J. Herrmann, Phys. Rev. Lett. **105** (2010).

[16] E. J. Friedman and A. S. Landsberg, Phys. Rev. Lett. **103**, 255701 (2009).

[17] H. D. Rozenfeld, L. K. Gallos, and H. A. Makse, Eur. Phys. J. B **75**, 305 (2010).

[18] F. Radicchi and S. Fortunato, Phys. Rev. E **81**, 036110 (2010).

[19] Y. S. Cho, B. Kahng, and D. Kim, Phys. Rev. E **81**, 030103(R) (2010).

[20] R. M. D'Souza and M. Mitzenmacher, Phys. Rev. Lett. **104** (2010).

[21] N. A. M. Araujo, J. S. Andrade, Jr., R. M. Ziff, and H. J. Herrmann, Phys. Rev. Lett. **106** (2011).

[22] S. Manna and A. Chatterjee, Physica A **390**, 177 (2011), ISSN 0378-4371.

[23] R. A. da Costa, S. N. Dorogovtsev, A. V. Goltsev, and J. F. F. Mendes, Phys. Rev. Lett. **105** (2010).

[24] P. Grassberger, C. Christensen, G. Bizhani, S.-W. Son, and M. Paczuski, Phys. Rev. Lett. **106** (2011).

[25] J. Nagler, A. Levina, and M. Timme, Nature Physics **7**, 265 (2011).

[26] O. Riordan and L. Warnke, arXiv (2011), 1102.5306.

[27] M. Plischke and B. Bergersen, *Equilibrium Statistical Physics, 2nd edition* (World Scientifc, Singapore, 1994).

[28] S. Boettcher et al., unpublished.

[29] S. N. Dorogovtsev, A. V. Goltsev, and J. F. F. Mendes, Rev. Mod. Phys. **80**, 1275 (2008).

[30] On a lattice, a fractal only exists in exactly one point, $p_l = p_u = p_c$.